

Intersecting delocalized p-branes

M. G. Ivanov*

July 24, 2001

Abstract

A model considered in the paper generalizes supergravity type model to the case of delocalized membrane sources. A generalization of intersecting p-brane solution with delocalized membranes is presented.

1 Introduction

Membrane theories [1, 2, 3, 4] consider membranes as mapping x from manifold $\mathbf{V}$ of dimension n to (pseudo)Riemannian manifold $\mathbf{M}$ of dimension $D > n$. The manifold $\mathbf{M}$ is interpreted as space-time (“bulk”). The image $x(\mathbf{V})$ is interpreted as membrane. Membrane appears in bulk equation in the form of singular sources localized at $x(\mathbf{V})$. A procedure of regularization can be considered as replacement of common membrane, which has zero thickness by some sort of thick membrane, which corresponds to continuous distribution of infinitely light membranes.

A covariant method of regularization of closed membrane theories (“ $(\mathbf{M} \rightarrow \mathbf{F})$ -approach”) was presented in the papers [10, 11]. $(\mathbf{M} \rightarrow \mathbf{F})$ -approach replaces mapping $x : \mathbf{V} \rightarrow \mathbf{M}$ by mapping $\varphi : \mathbf{M} \rightarrow \mathbf{F}$, where $\mathbf{F}$ is a manifold of dimension $D - n$. If for some point $\phi \in \mathbf{F}$ inverse image $\varphi^{-1}(\phi)$ is submanifold of dimension n , then it has to be considered as infinitely light membrane. The similar ideas were suggested before in papers [7, 8, 9] (see also references in paper [11]).

The generalization of p-brane solution with delocalized membranes was also presented in the papers [10, 11]. A natural generalization of extremal intersecting p-brane solution with delocalized membranes is constructed in the present paper (on intersecting p-brane solutions see [5, 6] and references in these papers).

2 Notation

D -dimensional space-time, i.e. smooth (pseudo)Riemannian manifold $\mathbf{M}$ with metric $ds^2 = g_{MN}dX^M dX^N$, $M, N, \dots = 0, \dots, D - 1$ is considered. In further calculations $\mathbf{M}$ is considered to be a finite region in $\mathbb{R}^D$ with smooth boundary $\partial\mathbf{M}$.

For differential forms A and B with components $A_{M_1 \dots M_q}$ and $B_{N_1 \dots N_p}$ the following tensor is introduced

$$(A, B)^{(k)}_{M_{k+1} \dots M_q N_{k+1} \dots N_p} = \frac{1}{k!} g^{M_1 N_1} \dots g^{M_k N_k} A_{M_1 \dots M_q} B_{N_1 \dots N_p}.$$

*e-mail: mgi@mi.ras.ru

Index (k) indicates the number of indices to contract. If it can not lead to ambiguity, (k) is skipped.

For differential form of power q it is convenient to introduce two norms $\|A\|_\Sigma^2 = \Sigma(A, A)^{(q)}$, where $\Sigma = \pm 1$, and Hodge duality operation $*A = (\Omega, A)^{(q)}$, where $\Omega = \sqrt{|g|} d^D X$ is form of volume. Here and below $g = \det(g_{MN})$, $\sigma = \text{sgn}(g)$.

3 Action and equations of motion

Let us consider the action of the following form

$$S = \int_{\mathbf{M}} d^D X \sqrt{|g|} \left(\frac{R}{2\kappa^2} - \frac{1}{2} \|\partial\phi\|^2 - \sum_I \frac{1}{2} \beta_I e^{-\alpha_I \cdot \phi} \|F^{(I)}\|^2 + \sum_a \lambda_{(I,a)} e^{\frac{1}{2} \alpha_I \cdot \phi} \|J^{(I,a)}\|_{\Sigma_{(I,a)} \sigma} + s_{(I,a)} \frac{\lambda_{(I,a)}}{h_{(I,a)}} (*J^{(I,a)}, A^{(I)}) \right). \quad (1)$$

Here $\phi = (\phi^1, \dots, \phi^{n_\phi})$ is a set of scalar fields (dilaton), $\alpha_I \cdot \phi = \sum_{i=1}^{n_\phi} \alpha_I^i \phi^i$, $F^{(I)} = dA^{(I)}$ are exact $(q_I + 1)$ -forms, $J^{(I,a)} = d\varphi_{(I,a)}^1 \wedge \dots \wedge d\varphi_{(I,a)}^{D-q_I}$, $\Sigma_{(I,a)} = \pm 1$, $s_{(I,a)} = \pm 1$, α_I^i , $\lambda_{(I,a)}$ and $h_{(I,a)}$ are real constants.

By variation of action (1) over metric g_{MN} , scalar fields ϕ , q_I -forms $A^{(I)}$ and membrane potentials $\varphi_{(I,a)}^\alpha$ one can find equations of motion. Einstein equations, found by variation of metric have the form

$$R_{MN} - \frac{1}{2} R g_{MN} = \kappa^2 T_{MN}, \quad (2)$$

where energy-momentum tensor is defined by the following formulae

$$T_{MN} = T_{MN}^{(\phi)} + \sum_I (T_{MN}^{(I)} + \sum_a T_{MN}^{(I,a)}), \quad (3)$$

$$T_{MN}^{(\phi)} = \partial_M \phi \cdot \partial_N \phi - \frac{1}{2} \|\partial\phi\|^2 g_{MN}, \quad (4)$$

$$T_{MN}^{(I)} = \beta_I e^{-\alpha_I \cdot \phi} \left((F^{(I)}, F^{(I)})_{MN} - \frac{1}{2} \|F^{(I)}\|^2 g_{MN} \right), \quad (5)$$

$$T_{MN}^{(I,a)} = -\lambda_{(I,a)} e^{\frac{1}{2} \alpha_I \cdot \phi} \|J^{(I,a)}\|_{\Sigma_{(I,a)} \sigma} \left(g_{MN} - \frac{(J^{(I,a)}, J^{(I,a)})_{MN}}{\|J^{(I,a)}\|^2} \right). \quad (6)$$

Equations, found by variation of forms $A^{(I)}$, have the form

$$\beta \delta \left(e^{-\alpha_I \cdot \phi} F^{(I)} \right) = \sigma (-1)^{q_I+1} Q^{(I)}, \quad (7)$$

where the source $Q^{(I)}$ defined by

$$Q^{(I)} = - \sum_a s_{(I,a)} \frac{\lambda_{(I,a)}}{h_{(I,a)}} * J^{(I,a)}. \quad (8)$$

Equations, found by variation of scalar fields, have the form

$$\square \phi = - \sum_I (Q_{\phi I} + Q_{\phi I s}), \quad (9)$$

where the sources are defined by

$$Q_{\phi I} = \frac{1}{2} \alpha_I \beta_I e^{-\alpha_I \cdot \phi} \|F^{(I)}\|^2, \quad (10)$$

$$Q_{\phi I s} = -\frac{\alpha_I}{2} e^{\frac{1}{2} \alpha_I \cdot \phi} \sum_a \lambda_{(I,a)} \|J^{(I,a)}\|_{\Sigma_{(I,a)} \sigma}. \quad (11)$$

Equations, found by variation of membrane potentials, have the form

$$\left(\delta \left[\Sigma_{(I,a)} \sigma \frac{e^{-\alpha_I \cdot \phi} J^{(I,a)}}{\|J^{(I,a)}\|_{\Sigma_{(I,a)} \sigma}} + (-1)^{q_I(D-q_I)} s_{(I,a)} \frac{*A^{(I,a)}}{h_{(I,a)}} \right], d\varphi_{(I,a)}^{\alpha_1} \wedge \dots \wedge d\varphi_{(I,a)}^{\alpha_{D-q_I-1}} \right) = 0. \quad (12)$$

4 Delocalized p-brane solutions

The goal of the paper is to present special solutions of equations (2), (7), (9) and (12).

One can introduce a set of orthonormal projectors $P_N^{(I,a) M}$, $\bar{P}_N^{(I,a) M} = \delta_N^M - P_N^{(I,a) M}$ numerated by index (I, a) . In the given coordinate system

$$\text{tr} \bar{P}^{(I,a)} = \bar{P}_M^{(I,a) M} = q_I; \quad P_N^{(I,a) M} = 0, \quad M \neq N, \quad (13)$$

i.e. $\bar{P}_N^{(I,a) M} = \Delta_{(I,a) M} \delta_{MN}$ (there is no summation over M), $\Delta_{(I,a) M} \in \{0, 1\}$.

Let us define fields by the following equations

$$A^{(I)} = \sum_a \frac{h_{(I,a)}}{H_{(I,a)}} \omega^{(I,a)}, \quad (14)$$

where

$$\omega^{(I,a)} = e^{W_{(I,a)}} \bigwedge_{\Delta_{(I,a) M}=1} dX^M, \quad (15)$$

$h_{(I,a)}$ are constants, $H_{(I,a)}$ are smooth positive functions (*parametrizing functions*), $W_{(I,a)}$ are smooth functions, which satisfy conditions $(P^{(I,a)} \partial)_M W_{(I,a)} = 0$.

$$ds^2 = \left(\prod_{I,a} H_{(I,a)}^{2\kappa^2 \Sigma_{(I,a)} \beta_I h_{(I,a)}^2 (\Delta_{(I,a) M} - \frac{q_I}{D-2})} \right) \eta_{MN} dX^M dX^N, \quad (16)$$

where $\eta_{MN} = \text{diag}(\pm 1, \dots, \pm 1)$.

$$\phi = \frac{1}{2} \sum_{I,a} \Sigma_{(I,a)} \beta_I h_{(I,a)}^2 \alpha_I \ln H_{(I,a)}. \quad (17)$$

$$J^{(I,a)} = \mathcal{J}_{(I,a)} * \omega^{(I,a)}. \quad (18)$$

Theorem on p -brane solutions with sources. *If fields defined by equations (14), (16), (17), (18), satisfy the following conditions*

$$|\mathcal{J}_{(I,a)}| = \frac{\Sigma_{(I,a)} \beta_I h_{(I,a)}^2}{\lambda_{(I,a)}} e^{-\frac{1}{2} \alpha_I \cdot \phi} \frac{\square H_{(I,a)}}{H_{(I,a)}^2}, \quad (19)$$

$$\text{sgn} \|\omega^{(I,a)}\|^2 = \Sigma_{(I,a)}, \quad (20)$$

$$(\bar{P}^{(I,a)} \partial)_M H_{(I,a)} = 0, \quad (21)$$

$$(P^{(I,a)}\partial)_M H_{(I,a)} (\bar{P}^{(I,a)}\partial)_N \left(\kappa^2 \sum_{J,b} \Sigma_{(J,b)} \beta_J h_{(J,b)}^2 \ln H_{(J,b)} 2\Delta_{(I,a)M} + W_{(I,a)} \right) = 0, \quad (22)$$

$$\ln H_{(I,a)} = \kappa^2 \sum_{J,b} \Sigma_{(J,b)} \beta_J h_{(J,b)}^2 \ln H_{(J,b)} \left[\frac{q_I q_J}{D-2} - \text{tr}(\bar{P}^{(I,a)} \bar{P}^{(J,b)}) - \frac{\alpha_I \cdot \alpha_J}{4\kappa^2} \right] + W_{(I,a)}, \quad (23)$$

$$\text{tr}(\bar{P}^{(I,b)} \bar{P}^{(I,c)}) \neq q_I - 1, \quad (24)$$

$$s_{(I,a)} = -\Sigma_{(I,a)} \sigma(-1)^{q_I(D-q_I)} \text{sgn} \mathcal{J}_{(I,a)}, \quad (25)$$

Then fields (14), (16), (17), (18) satisfy equations of motion (2), (7), (9), (12).

If $\square H_{(I,a)} = 0$ for all (I,a) , then membrane fields $J^{(I,a)}$ vanish and the solution is a regular intersecting extremal p-brane solution (see [5, 6] and references in these papers for examples). Vice-versa, one can introduce non-trivial membranes fields to intersecting extremal electric-type p-brane solution by setting $\square H_{(I,a)} \neq 0$.

5 Examples

As a simplest example, let us consider solution of Einstein-Maxwell equations in the presence dust cloud (with zero pressure) with charge density equal to mass density.

The action has the following form

$$S = \int d^4 X \sqrt{-g} \left(\frac{R}{2} - \frac{\|F\|^2}{2} - \|J\| - \frac{1}{\sqrt{2}} (*J, A) \right),$$

where $F = dA$ is electromagnetic field, A is 4-vector potential, $J = d\varphi^1 \wedge d\varphi^2 \wedge d\varphi^3$.

By variation over fields g_{MN} , A_M , φ^α one finds the following equations of motion

$$R_{MN} - \frac{1}{2} R g_{MN} = (F, F)_{MN} + \frac{(J, J)_{MN}}{\|J\|} - \left(\frac{1}{2} \|F\|^2 + \|J\| \right) g_{MN},$$

$$\left(\delta \left(\frac{J}{\|J\|} - \frac{*A}{\sqrt{2}} \right), d\varphi^\alpha \wedge d\varphi^\beta \right) = 0, \quad \delta F = -\frac{*J}{\sqrt{2}}.$$

The fields defined by the following equations solve the equations of motion

$$ds^2 = -H^{-2} dX^0 dX^0 + H^2 \delta_{\alpha\beta} dX^\alpha dX^\beta,$$

$$A = \frac{\sqrt{2}}{H} dX^0, \quad J = -2\Delta H dX^1 \wedge dX^2 \wedge dX^3.$$

Here H is smooth positive function of space coordinates X^α , $\alpha, \beta = 1, 2, 3$, $\Delta = \partial_1^2 + \partial_2^2 + \partial_3^2$. Calculating $\|J\|$ one has to choose a branch of square root, which gives $\|J\|\Delta H < 0$.

The following example describes intersecting delocalized 2-branes in 11d supergravity.

The action has the following form

$$S = \int d^{11} X \sqrt{-g} \left(\frac{R}{2} - \frac{\|F\|^2}{2} - \|J^{(1)}\| - \|J^{(2)}\| + \sqrt{2} (*J^{(1)} + *J^{(2)}, A) \right),$$

where $F = dA$, A is form of power 3, $J^{(a)} = d\varphi_a^1 \wedge \dots \wedge d\varphi_a^8$, $a = 1, 2$.

By variation over fields g_{MN} , A_{MNK} , φ_a^α one finds the following equations of motion

$$R_{MN} - \frac{1}{2}Rg_{MN} = (F, F)_{MN} + \frac{(J^{(1)}, J^{(1)})_{MN}}{\|J^{(1)}\|} + \frac{(J^{(2)}, J^{(2)})_{MN}}{\|J^{(2)}\|} + \left(\frac{1}{2}\|F\|^2 + \|J^{(1)}\| + \|J^{(2)}\| \right) g_{MN},$$

$$\left(\delta \left(\frac{J^{(a)}}{\|J^{(a)}\|} - \sqrt{2} * A \right), d\varphi_a^{\alpha_1} \wedge \dots \wedge d\varphi_a^{\alpha_7} \right) = 0, \quad \delta F = -\sqrt{2}(*J^{(1)} + *J^{(2)}).$$

The fields defined by the following equations solve the equations of motion

$$ds^2 = -H_1^{-2/3}H_2^{-2/3}dX^0dX^0 + H_1^{-2/3}H_2^{1/3}(dX^1dX^1 + dX^2dX^2)$$

$$+ H_1^{1/3}H_2^{-2/3}(dX^3dX^3 + dX^4dX^4) + H_1^{1/3}H_2^{1/3} \sum_{\alpha=5}^{10} dX^\alpha dX^\alpha,$$

$$A = \frac{1}{\sqrt{2}H_1} dX^0 \wedge dX^1 \wedge dX^2 + \frac{1}{\sqrt{2}H_2} dX^0 \wedge dX^3 \wedge dX^4,$$

$$J^{(1)} = -\frac{1}{2}\Delta H_1 dX^3 \wedge \dots \wedge dX^{10},$$

$$J^{(2)} = -\frac{1}{2}\Delta H_2 dX^1 \wedge dX^2 \wedge dX^5 \wedge dX^6 \wedge \dots \wedge dX^{10}.$$

Here H_a are smooth positive function of space coordinates X^α , $\alpha, \beta = 5, 6, 7, 8, 9, 10$, $\Delta = \sum_{\alpha=5}^{10} \partial_\alpha^2$. Calculating $\|J^{(a)}\|$ one has to choose a branch of square root, which gives $\|J^{(a)}\|\Delta H_a < 0$.

The author is grateful to I.V. Volovich and B. Dragovich. The work was partially supported by grant RFFI 99-01-00866.

References

- [1] M. Green, J.H. Schwarz, E. Witten, Superstring theory, vol.1,2, Cambridge University Press, 1987
- [2] B.M. Barbashov, V.V. Nesterenko, Model' relyativistkoj struny v fizike adronov, Moscow.: Energoatomizdat, 1987. 176 p. (in Russian)
- [3] V.S. Vladimirov, I.V. Volovich// Doklady Akademii Nauk. 1985. vol. 289. pp. 1043–1047
- [4] M.J. Duff, J.X. Lu// Nucl.Phys. B416, 1994. pp. 301–334; hep-th/9306052
- [5] I.Ya. Aref'eva, M.G. Ivanov, O.A. Rytchkov// Wess J., Akulov V.P.(Eds.), Supersymmetry and Quantum Field theory, Proceedings of the D. Volkov Memorial Seminar Held in Kharkov, Ukraine, 5–7 January 1997, Springer, pp. 25–41; hep-th/9702077
- [6] V.D. Ivashchuk, V.N. Melnikov// P-brane black holes for general intersections, Grav.Cosmol. 5, 1999. pp. 313–318; gr-qc/0002085
- [7] Y. Hosotani, Local scalar fields equivalent to Nambu-Goto strings, Phys.Rev.Letters 47 (1981) 399

- [8] T.R. Morris, From First to Second Quantized string Theory, Phys.Lett. 202B (1988) 222
- [9] L.M. Baker and D.B. Fairlie, Companion Equations for Branes, Journal of Mathematical Physics 41 (2000) 4284–4292; hep-th/9908157
- [10] M.G. Ivanov// The model of delocalized membranes, Doklady Akademii Nauk. 2001. vol. 378. pp. 26–28
- [11] M.G. Ivanov, Delocalized membrane model, hep-th/0105067